

**HIGHLIGHTS OF PRESCRIBING INFORMATION**

These highlights do not include all the information needed to use BAT safely and effectively. See full prescribing information for BAT.

**BAT, Botulism Antitoxin Heptavalent (A, B, C, D, E, F, G) - (Equine) Sterile Solution for Injection**  
**Initial U.S. Approval: 2013**

**INDICATIONS AND USAGE**

BAT [Botulism Antitoxin Heptavalent (A, B, C, D, E, F, G) – (Equine)] is a mixture of immune globulin fragments indicated for the treatment of symptomatic botulism following documented or suspected exposure to botulinum neurotoxin serotypes A, B, C, D, E, F, or G in adults and pediatric patients.

The effectiveness of BAT is based solely on efficacy studies conducted in animal models of botulism.

**DOSAGE AND ADMINISTRATION**

For intravenous use only.

Administer BAT by slow intravenous infusion after dilution 1:10 in normal saline at the dose recommended in the following table.

Patient Group	Dose	Starting Infusion Rate (first 30 minutes)	Incremental Infusion Rate if Tolerated (every 30 minutes)	Maximum Infusion Rate
Adults (≥ 17 years)	One vial	0.5 mL/min	Double the rate	2 mL/min
Pediatric (1 year to <17 years)	20 – 100% of adult dose	0.01 mL/kg/min Do not exceed the adult rate	0.01 mL/kg/min	0.03 mL/kg/min Do not exceed the adult rate
Infants (< 1 year)	10% of adult dose regardless of body weight	0.01 mL/kg/min	0.01 mL/kg/min	0.03 mL/kg/min

**DOSAGE FORMS AND STRENGTHS**

Each single-use vial contains a minimum potency of:

- 4,500 Units (U) for serotype A antitoxin
- 3,300 U for serotype B antitoxin,
- 3,000 U for serotype C antitoxin,
- 600 U for serotype D antitoxin,
- 5,100 U for serotype E antitoxin,
- 3,000 U for serotype F antitoxin, and
- 600 U for serotype G antitoxin

**CONTRAINDICATIONS**

None.

**WARNINGS AND PRECAUTIONS**

- Hypersensitivity reactions including anaphylaxis. Prepare for monitoring and management of allergic reactions (5.1).
- Delayed allergic reactions (serum sickness). Patient monitoring is recommended (5.2).
- Infusion reactions. Monitor and slow or interrupt infusion and administer treatment based on the severity of the reaction (5.3).
- Interference with non-glucose specific blood sugar testing systems. Use glucose-specific testing systems (5.4).
- BAT is made from equine plasma and may contain infectious agents e.g. viruses (5.5).

**ADVERSE REACTIONS**

- The most common adverse reactions observed in ≥ 5 % of healthy volunteers in clinical trials were headache, nausea, pruritus and urticaria (6.1).
- The most common adverse reactions reported in ≥ 1% of patients in a clinical study were pyrexia, rash, chills, nausea and edema (6.1).
- One serious adverse reaction of hemodynamic instability was observed in one patient in the clinical study (6.1).

To report SUSPECTED ADVERSE REACTIONS, contact Cangene Corporation at 1-800-768-2304 or FDA at 1-800-FDA-1088 or [www.fda.gov/medwatch](http://www.fda.gov/medwatch).

**USE IN SPECIFIC POPULATIONS**

- Pregnancy: No human or animal data. Use only if clearly needed (8.1).
- Pediatric: Limited safety data is available in the pediatric population. Dosing in pediatric patients is based on Salisbury Rule (8.4).

See 17 for PATIENT COUNSELING INFORMATION and FDA-approved patient labeling

Revised: 03/2013

**FULL PRESCRIBING INFORMATION: CONTENTS\***

1 INDICATIONS AND USAGE

2 DOSAGE AND ADMINISTRATION

    2.1 Dosage and Administration

    2.2 Preparation

3 DOSAGE FORMS AND STRENGTHS

4 CONTRAINDICATIONS

5 WARNINGS AND PRECAUTIONS

    5.1 Hypersensitivity Reactions

    5.2 Delayed Allergic Reactions (Serum Sickness)

    5.3 Infusion Reactions

    5.4 Interference with Blood Glucose Testing

    5.5 Transmissible Infectious Agents

6 ADVERSE REACTIONS

    6.1 Clinical Trials Experience

    6.2 Immunogenicity

7 DRUG INTERACTIONS

8 USE IN SPECIFIC POPULATIONS

    8.1 Pregnancy

    8.3 Nursing Mothers

    8.4 Pediatric Use

    8.5 Geriatric Use

11 DESCRIPTION

12 CLINICAL PHARMACOLOGY

    12.1 Mechanism of Action

    12.2 Pharmacodynamics

12.3 Pharmacokinetics

13 NONCLINICAL TOXICOLOGY

    13.2 Animal Toxicology and Pharmacology

14 CLINICAL STUDIES

15 REFERENCES

16 HOW SUPPLIED/STORAGE AND HANDLING

    16.1 How Supplied

    16.2 Storage and Handling

17 PATIENT COUNSELING INFORMATION

\*Sections or subsections omitted from the full prescribing information are not listed

## FULL PRESCRIBING INFORMATION

### 1 INDICATIONS AND USAGE

BAT [Botulism Antitoxin Heptavalent (A, B, C, D, E, F, G) – (Equine)] is a mixture of immune globulin fragments indicated for the treatment of symptomatic botulism following documented or suspected exposure to botulinum neurotoxin serotypes A, B, C, D, E, F, or G in adults and pediatric patients.

The effectiveness of BAT is based on efficacy studies conducted in animal models of botulism.

### 2 DOSAGE AND ADMINISTRATION

**BAT is for intravenous use only.**

#### 2.1 Dosage and Administration

- Each vial of BAT contains a minimum potency for serotypes A, B, C, D, E, F, and G antitoxin [3 DOSAGE FORMS AND STRENGTHS].
- For adult, pediatric, and infant patient groups, administer a dose of BAT according to Table 1. For details on pediatric dosing by body weight see Table 2.
- Consider skin sensitivity testing for patients at risk of acute hypersensitivity reactions [5.1 Hypersensitivity Reactions].

##### *Skin Sensitivity Test*

Administer 0.02 milliliters of 1:1000 saline-diluted BAT (enough to raise a small wheal) intradermally on the volar surface of the forearm. If the test is negative, repeat the test using a 1:100 dilution. Perform concurrent positive (histamine) and negative (saline) control tests. A positive test is a wheal with surrounding erythema at least 3 millimeters larger than the negative control test; read at 15-20 minutes. The histamine control must be positive for valid interpretation.

- Administer all BAT doses after dilution 1:10 in normal saline by slow intravenous infusion according to the varying infusion rates in Table 1.
- Monitor vital signs throughout the infusion. If tolerated, the infusion rate can be increased incrementally up to the maximum infusion rate, and continued for the remainder of the administration. Decrease infusion rate if the patient develops discomfort or infusion-related adverse reactions.

**Table 1 BAT Dosing Guide and Intravenous Infusion Rate**

Patient Group	Dose	Starting Infusion Rate (first 30 minutes)	Incremental Infusion Rate if Tolerated (every 30 minutes)	Maximum Infusion Rate
Adults (≥ 17 years)	One vial	0.5 mL/min	Double the rate	2 mL/min
Pediatric (1 year to <17 years)	20 – 100% of adult dose	0.01 mL/kg/min Do not exceed the adult rate.	0.01 mL/kg/min	0.03 mL/kg/min Do not exceed the adult rate
Infants (< 1 year)	10% of adult dose regardless of body weight	0.01 mL/kg/min	0.01 mL/kg/min	0.03 mL/kg/min

Calculate pediatric BAT dose by body weight according to Table 2.

**Table 2 Pediatric Dosing Guide for BAT Based on Salisbury Rule**

Body Weight (kg)	Percent of Adult Dose* (%)
10-14	20**
15-19	30
20-24	40
25-29	50
30-34	60
35-39	65
40-44	70
45-49	75
50-54	80
≥ 55	100

\*Dosing guide is based on the Salisbury Rule (1):

- Body weight ≤ 30 kg:  $2 \times \text{weight (kg)} = \% \text{ adult dose to administer}$
- Body weight > 30 kg:  $\text{weight (kg)} + 30 = \% \text{ adult dose to administer}$

Do not exceed 1 vial dose regardless of body weight.

\*\* Minimum dose is 20% of adult dose.

## 2.2 Preparation

1. Bring vial to room temperature.

- If frozen, thaw vial by placing in a refrigerator at 36 to 46 °F (2 to 8 °C) until the contents are thawed for approximately 14 hours.

- Product can be thawed rapidly by placing at room temperature for one hour followed by a water bath at 98.6 °F (37 °C) until thawed.
  - Do not thaw this product in a microwave oven. Do not refreeze the vial.
2. Inspect vial to ensure there is no damage to the seal or vial. If damaged, discard the vial.
  3. Do not shake the vial during preparation to avoid foaming.
  4. Dilute 1:10 in 0.9% Sodium Chloride Injection, USP (saline) by adding BAT solution from the vial to the appropriate amount of saline in an IV bag. Do not use any other diluents. As the fill volume per vial varies by lot number (approximately 10 to 22 milliliters per vial), 90 to 200 milliliters of saline will be required. Withdraw the entire contents of the vial to obtain the total volume in the vial. If a partial vial is required (for pediatric dosing), the entire content of the vial should be withdrawn to ensure accurate calculation of the dosage [Table 2].
  5. Visually inspect the product for particulate matter and discoloration prior to administration. Do not use if the solution is turbid, cloudy, or contains particles.
  6. Use an intravenous line with constant infusion pump. Use of an in line filter is optional.
  7. BAT vials are for single use only and contain no preservative.
  8. Discard any unused portion.

### **3 DOSAGE FORMS AND STRENGTHS**

BAT is a sterile solution of purified F(ab')<sub>2</sub> plus F(ab')<sub>2</sub>-related immune globulin fragments derived from equine plasma, containing antitoxin activity to botulinum neurotoxins A, B, C, D, E, F, and G.

Each single-use vial, regardless of size or fill volume, contains a minimum antitoxin potency of:

- 4,500 U serotype A antitoxin,
- 3,300 U serotype B antitoxin,
- 3,000 U serotype C antitoxin,
- 600 U serotype D antitoxin,
- 5,100 U serotype E antitoxin,
- 3,000 U serotype F antitoxin, and
- 600 U serotype G antitoxin.

### **4 CONTRAINDICATIONS**

None.

## 5 WARNINGS AND PRECAUTIONS

### 5.1 Hypersensitivity Reactions

Severe hypersensitivity reactions, including anaphylactic and anaphylactoid reactions may occur following BAT administration. Patients with a history of hypersensitivity to horses or equine blood products, asthma, and hay fever are at a greater risk for developing severe hypersensitivity reactions to BAT. To ascertain risk of allergic reactions in these cases, consider performing a skin sensitivity test [see 2.1 Dosage and Administration].

Administer BAT in a setting with appropriate equipment, medication, and personnel trained in the management of hypersensitivity, anaphylaxis, and shock.

Monitor all patients for signs and symptoms of acute allergic reaction (e.g. urticaria, pruritus, erythema, angioedema, bronchospasm with wheezing or cough, stridor, laryngeal edema, hypotension, tachycardia) during and following the BAT infusion. In case of hypersensitivity reaction, discontinue BAT administration immediately and administer appropriate emergency care. Medications such as epinephrine should be available for immediate treatment of acute hypersensitivity reactions.

For patients at risk for hypersensitivity reaction, begin BAT administration at the lowest rate achievable (<0.01 mL/min) and monitor.

### 5.2 Delayed Allergic Reactions (Serum Sickness)

Delayed allergic reactions (serum sickness e.g. fever, urticarial or maculopapular rash, myalgia, arthralgia, and lymphadenopathy) may occur following BAT administration, typically 10-21 days after infusion. Monitor patients for signs and symptoms of delayed allergic reaction.

If a delayed allergic reaction (serum sickness) is suspected, administer appropriate medical care.

### 5.3 Infusion Reactions

Chills, fever, headaches, nausea, and vomiting can be related to the rate of infusion. Arthralgia, myalgia and fatigue or vasovagal reactions may also develop. Carefully observe patients for the onset of these infusion reactions throughout the infusion period and immediately following an infusion.

Reduce the rate of infusion if the patient experiences infusion reactions and administer symptomatic therapy. If symptoms worsen, discontinue the infusion and administer appropriate medical care.

### 5.4 Interference with Blood Glucose Testing

The maltose contained in BAT can interfere with some types of blood glucose monitoring systems i.e. those based on glucose dehydrogenase pyrroloquinoline-quinone (GDH-PQQ) method. This can result in falsely elevated glucose readings and inappropriate administration of insulin, resulting in life-threatening hypoglycemia. Cases of true hypoglycemia may go

untreated if the hypoglycemic state is masked by falsely elevated results [7 DRUG INTERACTIONS].

## 5.5 Transmissible Infectious Agents

Because BAT is made from equine plasma, it may carry the risk of transmitting infectious agents e.g. viruses. The equine plasma pools are screened for the presence of certain infectious agents and the manufacturing process for BAT includes measures to inactivate and remove certain viruses [11 DESCRIPTION]. Despite these measures, such products can still potentially transmit disease. No cases of transmission of viral diseases have been associated with the use of BAT.

Report all infections thought by a physician to have been transmitted by BAT to Cangene Corporation at 1-800-768-2304. Discuss the risks and benefits of this product with the patient or their legal guardian before administering it to the patient [17 PATIENT COUNSELLING INFORMATION].

## 6 ADVERSE REACTIONS

The most common adverse reactions observed in  $\geq 5\%$  of healthy volunteers in clinical trials were headache, nausea, pruritus, and urticaria.

The most common adverse reactions reported in  $\geq 1\%$  of patients in a clinical study were pyrexia, rash, chills, nausea and edema.

The following serious adverse reactions are discussed in detail in other sections of the labeling:

- Hypersensitivity reactions [5.1]
- Delayed allergic reactions/serum sickness [5.2]
- Infusion reactions [5.3]

### 6.1 Clinical Trials Experience

*Because clinical trials are conducted under widely varying conditions, adverse reaction rates observed in the clinical trials of a drug cannot be directly compared to rates in the clinical trials of another drug and may not reflect the rates observed in practice.*

In a randomized, double-blind, parallel arm trial conducted to evaluate the safety of BAT in healthy subjects, and to establish the pharmacokinetic profile of the seven botulinum antitoxin serotypes contained in BAT following intravenous (IV) administration, 40 subjects were randomized to receive either one (n=20) or two vials (n=20) of BAT.

In a second parallel arm, randomized, double-blind pharmacodynamic trial, 26 healthy subjects were randomized to receive either BAT in saline (n=16) or placebo (0.9% saline; n=10).

The most common adverse reactions in all healthy subjects were headache (9%), pruritus (5%), nausea (5%), and urticaria (5%). Other adverse reactions reported in less than 4% of subjects included pyrexia and throat discomfort. All reported adverse reactions were

considered mild or moderate. No serious adverse reactions were reported. Two moderate acute allergic reactions that required premature termination of the infusion and treatment were reported. Reactions were predefined as mild if the subject was aware but could tolerate. Moderate reactions were predefined as discomfort enough to interfere with normal daily activity.

A total of 231 subjects with suspected or confirmed botulism were exposed to BAT in an open-label observational expanded access clinical study sponsored by the Centers for Disease Control and Prevention (CDC).

The majority of adult (213/216) and pediatric (13/15) subjects received one dose of BAT. Three adult subjects were exposed to a second dose of BAT, and two pediatric subjects each received two infant doses (10% of the adult dose). The administration of a second dose varied from seven hours to one month after the first dose.

Safety data was actively collected from treating physicians by the CDC. However, no on-site safety monitoring was performed, and the CDC relied on follow-up information provided by the treating physicians to determine the reporting frequencies for adverse reactions. Of the 231 subjects receiving BAT, safety information was available for 228 subjects. Adverse reactions were reported in 10% of all subjects. The most common adverse reactions were pyrexia (4%), rash (2%), chills (1%), nausea (1%), and edema (1%). Other adverse reactions were reported in less than 1% of subjects. No subject experienced anaphylaxis. One subject experienced a serious adverse reaction of hemodynamic instability characterized by bradycardia, tachycardia, and asystole during BAT administration. One subject experienced mild serum sickness (<1%) with myalgia, arthralgia, and dark urine twelve days after BAT administration.

**Table 3 Summary of Adverse Drug Reactions (ADR) Reported in Subjects that Received BAT through the CDC Expanded Access Clinical Study**

System Organ Class	Preferred Term	Overall (N=228)		
		No. of Events	No. of Subjects	% of Subjects
ALL BODY SYSTEM	OVERALL	37	23	10.1
Cardiac disorders	Cardiac arrest	1	1	0.4
	Bradycardia	1	1	0.4
	Tachycardia	1	1	0.4
Gastrointestinal disorders	Vomiting	1	1	0.4
	Nausea	2	2	0.9
General disorders and administration site conditions	Pyrexia	9	9	3.9
	Chest discomfort	1	1	0.4
	Edema	2	2	0.9
	Chills	3	3	1.3
	Feeling jittery	1	1	0.4
Immune system disorders	Serum Sickness	1	1	0.4
Investigations	Blood pressure increased	1	1	0.4

System Organ Class	Preferred Term	Overall (N=228)		
		No. of Events	No. of Subjects	% of Subjects
	White blood cell count increased	1	1	0.4
Psychiatric disorders	Agitation	1	1	0.4
	Anxiety	1	1	0.4
Renal and urinary disorders	Urinary retention	1	1	0.4
Respiratory, thoracic and mediastinal disorders	Bronchospasm	1	1	0.4
Skin and subcutaneous tissue disorders	Erythema	1	1	0.4
	Hyperhidrosis	1	1	0.4
	Rash	4	4	1.8
Vascular disorders	Hemodynamic instability	1	1	0.4
	Hypotension	1	1	0.4

All adverse reactions were classified according to MedDRA Version 15.0 and are ranked according to medical significance within a given SOC.

## 6.2 Immunogenicity

As with all therapeutic proteins, there is potential for immunogenicity. All subjects from the two clinical trials were tested for immunogenicity against BAT at baseline and at the end of the studies (Day 28) using a validated assay. Eleven subjects seroconverted during the course of the two trials. One subject from each clinical trial experienced a moderate allergic reaction during the administration of BAT. Both subjects were negative for anti-BAT antibodies at baseline and at the end of their respective studies. The detection of antibody formation is highly dependent on the sensitivity and specificity of the assay. Additionally, the observed incidence of antibody (including neutralizing antibody) positivity in an assay may be influenced by several factors including assay methodology, sample handling, timing of sample collection, concomitant medications, and underlying disease. For these reasons, comparison of the incidence of antibodies to BAT with the incidence of antibodies to other products may be misleading.

## 7 DRUG INTERACTIONS

### *Drug Laboratory Interactions: Blood Glucose Testing*

BAT contains maltose which can interfere with certain types of blood glucose monitoring systems [5.4 Interference with Blood Glucose Testing]. Only test systems that are glucose-specific should be used in patients receiving BAT. This interference can result in falsely elevated glucose readings that can lead to untreated hypoglycemia or to inappropriate insulin administration, resulting in life-threatening hypoglycemia.

The product information of the blood glucose testing system, including that of the test strips, should be carefully reviewed to determine if the system is appropriate for use with maltose-containing parenteral systems. If any uncertainty exists, contact the manufacturer of the



testing system to determine if the system is appropriate for use with maltose-containing parenteral products.

## **8 USE IN SPECIFIC POPULATIONS**

### **8.1 Pregnancy**

Pregnancy Category C. Animal reproduction studies have not been conducted with BAT. It is not known whether BAT can cause fetal harm when administered to a pregnant woman or can affect reproductive capacity. BAT should only be given to pregnant women if the benefits outweigh the risks.

### **8.3 Nursing Mothers**

It is not known whether BAT is excreted in human milk. Caution should be exercised when BAT is administered to a nursing mother since many drugs are excreted in human milk.

### **8.4 Pediatric Use**

The effectiveness of BAT has not been established in pediatric patients. Limited pediatric safety data are available.

Fifteen pediatric subjects (age 10 days to 17 years; including 1 newborn, 3 infants and toddlers, 4 children and 7 adolescents) received BAT under the CDC expanded access clinical study. A 3-year old subject and an infant received two infant doses, and 13 pediatric subjects received one pediatric dose according to Salisbury Rule [Table 2].

Two adverse reactions were reported in two pediatric subjects. One subject experienced an adverse reaction of pyrexia following infusion of BAT, while the other subject experienced a serious adverse reaction of hemodynamic instability characterized by tachycardia, bradycardia, and asystole during infusion of BAT.

Dosing in pediatric patients is based on Salisbury Rule.

### **8.5 Geriatric Use**

The safety, pharmacokinetics, and effectiveness of BAT have not been established in geriatric subjects.

Thirty six geriatric subjects received BAT under the CDC expanded access clinical study. One geriatric subject experienced rash as an adverse reaction following infusion of BAT.

## **11 DESCRIPTION**

BAT [Botulism Antitoxin Heptavalent (A, B, C, D, E, F, G) – (Equine)] is a sterile solution of F(ab')<sub>2</sub> and F(ab')<sub>2</sub>-related antibody fragments prepared from plasma obtained from horses that have been immunized with a specific serotype of botulinum toxoid and toxin. To obtain the final heptavalent product, the seven antitoxin serotypes are blended. BAT is supplied in

either a 20 or 50 milliliter vial size, with a fill volume ranging from 10 to 22 milliliters per vial. BAT is administered intravenously.

The manufacturing process for each antitoxin type includes cation-exchange chromatography to purify the immune globulin fraction, digestion with pepsin to produce F(ab')<sub>2</sub> plus F(ab')<sub>2</sub>-related immune globulin fragments, anion exchange chromatography to remove the pepsin as well as other impurities and filtration. In addition, the manufacturing process includes two viral inactivation/removal steps; solvent/detergent (S/D) treatment and virus filtration [Table 4].

The S/D treatment step using tri-n-butyl phosphate (TnBP) and triton X-100 (TX-100) is effective at inactivating known lipid-enveloped viruses such as equine encephalitis, equine arteritis, West Nile virus, equine infectious anemia, equine herpes virus, rabies, and equine influenza. The BAT manufacturing process also includes a robust filtration step that is effective in reducing the levels of some lipid-enveloped viruses (listed above) as well as non-enveloped viruses including equine rhinovirus, equine adeno- and adeno-associated viruses, and equine parvovirus.

**Table 4 Viral Clearance Capacity of the BAT Process**

Genome	Enveloped					Non-enveloped		
	RNA	RNA	RNA	DNA	RNA	DNA	RNA	RNA
Virus	XmuLV	WNV	BVDV	PRV	PI3	Ad2	EMC	Porcine Parvovirus
Family	Retro	Flavi	Flavi	Herpes	Paramyxo	Adeno	Picornia	Parvo
Size (nm)	80-110	40-70	50-70	150-200	100-200	70-90	25-30	18-24
Nanofiltration (log <sub>10</sub> )	≥ 2.7	≥ 2.1	≥ 4.5			≥ 4.7	≥ 4.5	4.5
S/D (log <sub>10</sub> )	≥ 4.3	≥ 5.1		≥ 5.1	≥ 5.5			
Total Reduction (log <sub>10</sub> )	≥ 7.0	≥ 7.2	≥ 4.5	≥ 5.1	≥ 5.5	≥ 4.7	≥ 4.5	4.5

XmuLV: Xenotropic Murine Leukemia Virus; specific model for equine infectious anemia, and a model for lipid-enveloped RNA viruses of similar size, such as vesicular stomatitis virus (Rhabdo family).

WNV: West Nile Virus; relevant virus, and specific model for lipid-enveloped RNA viruses, including the arboviruses, which contains both Flaviviridae and Togaviridae and includes equine encephalitis viruses (Toga family) and equine viral arteritis (Arteri family, formerly a Toga virus).

BVDV: Bovine Viral Diarrhea Virus; relevant virus, and specific model for lipid-enveloped RNA viruses, including the arboviruses, which contains both Flaviviridae and Togaviridae and includes equine encephalitis viruses (Toga family) and equine viral arteritis (Arteri family, formerly a Toga virus).

PRV: Pseudorabies Virus; specific model for equine herpes viruses and non-specific model for lipid-enveloped viruses.

PI3: Parainfluenza III Virus; model for lipid enveloped RNA viruses, and viruses of the similar family, orthomyxo, which includes equine influenza virus.

Ad2: Adenovirus; specific model for equine adenovirus.

EMC: Encephalomyocarditis Virus; specific model for equine parvovirus and adeno-associated virus, non-specific model for small lipid and non-lipid enveloped viruses.

BAT is formulated with 10% maltose and 0.03% polysorbate 80. The formulated bulk material contains approximately 3-7 g% (30-70 milligrams/milliliter) protein.

The product potency is expressed in units based on the mouse neutralization assay (MNA). Each unit of BAT is designed to neutralize 10,000 mouse intraperitoneal lethal dose 50%

units (MIPLD<sub>50</sub>) of botulinum neurotoxin for serotype A, B, C, D, F, and G and 1,000 MIPLD<sub>50</sub> of serotype E.

## **12 CLINICAL PHARMACOLOGY**

### **12.1 Mechanism of Action**

The mechanism of action of BAT is through passive immunization with equine polyclonal antibody fragments (primarily F(ab')<sub>2</sub> and Fab) against botulinum neurotoxin (BoNT) A, B, C, D, E, F, and G. In the circulation the polyclonal antibody fragments bind to free BoNT. This prevents the BoNT from interacting with ganglioside anchorage sites and protein receptors on the cholinergic nerve endings. In turn this prevents BoNT internalization into the target cells. The antibody/antigen complexes are then cleared from the circulation by the organs involved in processing immune complexes.

Experimental evidence concerning the amount of circulating antitoxin needed to counteract BoNT intoxication is not fully documented. The outcome of treatment depends, as it does with other comparable conditions, largely on the time interval elapsing after the onset of symptoms and antitoxin administration.

### **12.2 Pharmacodynamics**

A proof-of-concept clinical dose-response trial was conducted using the extensor digitorum brevis (EDB) muscle of the foot as a model for measuring muscle paralysis after exposure to botulinum toxin. In this model, BAT prevented subjects from experiencing a decrease in muscle function after exposure to botulinum neurotoxin (BoNT) serotypes A and B. Subjects treated with placebo (n=10) demonstrated a loss of greater than 50% EDB muscle function within 3 days of exposure to BoNT serotypes A and B. In the BAT arm of the trial (n=16), EDB muscle function was stable over time indicating that BAT was effective in preserving muscle function for up to 28 days following exposure to both BoNT serotype A and B.

### **12.3 Pharmacokinetics**

The pharmacokinetics (PK) of the seven botulinum antitoxin serotypes was determined in healthy human subjects following IV administration of either one (n=20) or two vials (n=20) of BAT. The various PK parameters are summarized in Table 5.

The PK parameters varied based upon the antitoxin serotype measured. Antitoxin serotypes D and E had the shortest half-lives. While antitoxin serotype B and C had the longest half-lives. The AUC<sub>0-∞</sub> and C<sub>max</sub> values increased in a dose proportional fashion as the BAT dose increased from one to two vials. In addition, mean clearance values appeared to be similar between both treatment groups for the seven antitoxin serotypes, suggesting dose linearity of BAT over the dose range studied.

**Table 5 Pharmacokinetic Parameters (Mean) for Antitoxin Serotypes A Through G in Humans Following Intravenous Administration of either One or Two Vials of BAT**

Antitoxin Serotype	Treatment Group	AUC <sub>0-∞</sub> (U*hr/mL)	C <sub>max</sub> (U/mL)	t <sub>1/2</sub> (hr)	Cl (mL/hr)	V <sub>d</sub> (mL)
A	1 Vial	26.00	2.69	8.64	293	3637
	2 Vials	56.09	6.23	10.20	285	3993
B	1 Vial	29.30	1.90	34.20	196	9607
	2 Vials	62.55	4.28	57.10	181	14865
C	1 Vial	37.34	2.26	29.60	144	6066
	2 Vials	86.25	4.89	45.60	127	8486
D	1 Vial	7.62	0.81	7.51	137	1465
	2 Vials	14.83	1.60	7.77	151	1653
E	1 Vial	7.16	0.94	7.75	1250	14172
	2 Vials	15.66	1.75	7.32	1110	11596
F	1 Vial	31.40	2.37	14.10	169	3413
	2 Vials	63.19	4.29	18.20	168	4334
G	1 Vial	7.05	0.59	11.70	149	2372
	2 Vials	14.66	1.19	14.70	144	3063

AUC = Area Under the Concentration Curve; Cl = Clearance; C<sub>max</sub> = Maximum Serum Concentration; BAT = Botulism Antitoxin Heptavalent (A, B, C, D, E, F, G) – (Equine); t<sub>1/2</sub> = Half-life; T<sub>max</sub> = Time to Maximum Serum Concentration; U = Unit; V<sub>d</sub> = Volume of Distribution.

## 13 NONCLINICAL TOXICOLOGY

### 13.2 Animal Toxicology and Pharmacology

Toxicological studies were not conducted for BAT or its components.

The evaluation of new treatment options for botulism using controlled human trials is unethical and infeasible. Therefore the effectiveness of BAT for treatment of botulism is based on well controlled efficacy studies conducted in guinea pigs and Rhesus macaques.

#### *Guinea Pig*

In a controlled therapeutic efficacy study, guinea pigs were intoxicated with various BoNT serotypes (A, B, C, D, E, F or G) at a dose of 1.5x guinea pig intramuscular lethal dose 50% units (GPIMLD<sub>50</sub>) via intramuscular injection into the right hind limb. The animals were then treated with either placebo control or 1x scaled human dose of BAT (weight/weight based on an average human body weight of 70 kilograms), after the onset of moderate clinical signs of botulism (right hind limb weakness, salivation, lacrimation, weak limbs and noticeable changes in breathing rate or pattern). Treatment with BAT resulted in a statistically significant improvement in the survival rate of animals across all of the serotypes tested [Table 6].

**Table 6 Summary of Guinea Pig Survival Data from BAT Therapeutic Efficacy Study**

Neurotoxin Serotype	Treatment Group	Survival Rate (%)	Two-sided Fisher's Exact Test (p-value)
A	1x BAT	34/34 (100%)	p<0.0001
	Placebo Control	0/34 (0%)	
B	1x BAT	34/34 (100%)	p<0.0001
	Placebo Control	1/34 (3%)	
C	1x BAT	33/34 (97%)	p<0.0001
	Placebo Control	4/34 (12%)	
D	1x BAT	33/34 (97%)	p<0.0001
	Placebo Control	5/34 (15%)	
E	1x BAT	34/34 (100%)	p<0.0001
	Placebo Control	0/34 (0%)	
F	1x BAT	34/34 (100%)	p<0.0001
	Placebo Control	4/34 (12%)	
G	1x BAT	34/34 (100%)	p<0.0001
	Placebo Control	17/34 (50%)	

BAT = Botulism Antitoxin Heptavalent (A, B, C, D, E, F, G) – (Equine).

#### *Non-human Primate*

In a controlled therapeutic efficacy study, rhesus macaques were intoxicated with BoNT serotype A delivered intravenously at a dose of 1.7x nonhuman primate intravenous lethal dose 50% (NHPLD<sub>50</sub>) units per kilogram of body weight. The animals were then treated with either placebo control or 1x scaled human dose of BAT (weight/weight based on an average human body weight of 70 kilograms), after the onset of clinical signs of botulism (ptosis, muscular weakness, or respiratory distress). Treatment with BAT resulted in a statistically significant improvement in the survival rate [Table 7].

**Table 7 Summary of Rhesus macaque Survival Data from BAT Therapeutic Efficacy Study**

Treatment Group	Survival Rate (%)	Two-sided Fisher's Exact Test (p-value)
1x BAT	14/30 (47%)	p<0.0001
Placebo Control	0/30 (0%)	

BAT = Botulism Antitoxin Heptavalent (A, B, C, D, E, F, G) – (Equine).

## 14 CLINICAL STUDIES

The effectiveness of BAT is based on efficacy studies demonstrating a survival benefit in animal models of botulism [13.2 Animal Toxicology and Pharmacology]. The safety has been tested in healthy adults and patients with suspected botulism who were treated with BAT under an expanded access clinical study.

The pharmacokinetic, pharmacodynamic, and safety profiles of BAT have been evaluated in two clinical studies. In these clinical studies, BAT was shown to have an acceptable safety profile when one or two vials of BAT were administered intravenously to healthy subjects.

In a randomized, single-center, double-blind trial the pharmacokinetics and safety of BAT was evaluated in 40 healthy subjects receiving either one (n = 20) or two (n = 20) vials of BAT by IV infusion. Serum BAT levels were measured in the subjects using the Mouse Neutralization Assay (MNA). A brief summary of the results can be found in [12.3 Pharmacokinetics].

In a randomized single center, double-blind trial the pharmacodynamics and safety of BAT was evaluated in 26 healthy subjects receiving either a single vial of BAT (n=16) or placebo (n=10) by IV infusion. The effects of BAT in preventing paralysis of the EDB foot muscle following administration of botulinum neurotoxin serotype A or B was determined. A brief summary of the results can be found in [12.2 Pharmacodynamics].

To provide additional support for the efficacy demonstrated in the animal models, a preliminary analysis of data from a Centers for Disease Control and Prevention (CDC) open-label, observational expanded access clinical study for the treatment of subjects with suspected or confirmed botulism with BAT was conducted. Across the 148 subjects treated with BAT in the period analyzed, 109 subjects had a final discharge diagnosis of suspected or confirmed botulism and were included in the analysis population. The median time from the onset of botulism symptoms to treatment with BAT was 3.6 days (range: 0.25 – 38 days). Early treatment ( $\leq 2$  days after onset of symptoms) with BAT was associated with a shorter length of hospitalization, duration in intensive care unit (ICU) and duration of mechanical ventilation compared to later treatment [Table 8] and is consistent with the mechanism of action [12.1 Mechanism of Action].

**Table 8 Summary of Duration of Hospitalization, ICU Stay and Mechanical Ventilation for CDC Patients Treated with BAT**

	Time from Symptoms to Treatment	Number of Patients (N)	Mean Duration in Days (SD)
Hospitalization	$\leq 2$ Days	14	12.4 (9.28)
	$> 2$ Days	72	26.1 (26.37)
ICU Stay	$\leq 2$ Days	13	9.2 (7.40)
	$> 2$ Days	70	15.8 (18.76)
Mechanical Ventilation	$\leq 2$ Days	9	11.6 (7.83)
	$> 2$ Days	41	23.4 (21.11)

## 15 REFERENCES

1. Lack JA, Stuart-Taylor ME. Calculation of drug dosage and body surface area of children. Br J Anaesth. 1997; 78:601-605.

## 16 HOW SUPPLIED/STORAGE AND HANDLING

### 16.1 How Supplied

BAT is supplied in either 20 milliliter or 50 milliliter glass vials sealed with a butyl rubber stopper and an aluminum seal with a plastic flip-top cap, with a fill volume ranging from 10 to 22 milliliters per vial. Each vial, regardless of size or fill volume contains a minimum potency of >4,500 U serotype A antitoxin, >3,300 U serotype B antitoxin, >3000 U serotype C antitoxin, >600 U serotype D antitoxin, >5,100 U serotype E antitoxin, >3,000 U serotype F antitoxin, and >600 U serotype G antitoxin.

BAT is not made with natural rubber latex.

<b>NDC Number</b>	<b>Product Description</b>
<b>60492-0075-2</b>	A 50 milliliter single dose vial.
<b>60492-0075-3</b>	A 20 milliliter single dose vial.

### 16.2 Storage and Handling

- Store frozen at or below  $\leq 5^{\circ}\text{F}$  ( $\leq -15^{\circ}\text{C}$ ) until used.
- Once thawed, Botulism Antitoxin Heptavalent (A, B, C, D, E, F, G) - (Equine) may be stored for a maximum of 36 months at 2-8 °C (36-48 °F) or until 48 months from the date of manufacture, whichever comes first. Do not refreeze.

## 17 PATIENT COUNSELING INFORMATION

See FDA-approved patient labeling (Patient Information)

- Inform patients of the following:
  - BAT is prepared from equine plasma and may contain infectious agents such as viruses that can cause disease.
  - The risk that such products will transmit an infectious agent has been reduced by screening the horses for prior exposure to certain viruses, by testing for the presence of certain current viral infections, and by inactivating and/or removing certain viruses during manufacturing.
  - Despite these measures, such products can still potentially transmit disease.
  - There is also the possibility that unknown infectious agents may be present in such products.
- Inform patients that persons known to have allergies to horses or other allergies such as seasonal allergies or hay fever, or asthma may be at increased risk of hypersensitivity reactions and should only receive BAT if the benefits outweigh the risks.
- Advise patients about the potential interference with non-glucose specific monitoring systems.

- The maltose contained in BAT can interfere with some types of blood glucose monitoring systems.
- Only testing systems that are glucose-specific should be used in patients receiving BAT.
- This interference can result in falsely elevated glucose readings that can lead to untreated hypoglycemia or to inappropriate insulin administration, resulting in life-threatening hypoglycemia.



## PATIENT INFORMATION

BAT [Botulism Antitoxin Heptavalent (A, B, C, D, E, F, G) – (Equine)]

### **What is botulism?**

Botulism is a muscle-paralyzing disease caused by a toxin made by a bacterium called *Clostridium botulinum*.

Botulism can cause the following conditions:

- Double vision,
- Blurred vision,
- Drooping eyelids,
- Slurred speech,
- Difficulty swallowing,
- Dry mouth,
- Muscle weakness that spreads through the body,
- Difficulty breathing.

Botulism can also cause paralysis and death. After a person is exposed to the toxin, problems can start as early as three hours or as late as a few days. It can take weeks or months to get better. During that time, many people need special care in the hospital.

The effectiveness of BAT has been studied in animals with botulism.

### **What is BAT?**

BAT is a botulism antitoxin made from the plasma of horses. It contains antibody fragments which can neutralize botulism toxins. BAT may make the illness from botulism less severe. Treatment with BAT will not reverse the paralysis, but may decrease the duration and extent of paralysis.

### **Who should use BAT?**

Your doctor may give you BAT if they suspect that you have been exposed to botulism toxin. You should get the treatment as quickly as possible to stop the progression of the illness.

Unless the benefits outweigh the risks, you should not receive BAT if you have a known history of allergies to horses or horse blood products.

### **How will you receive BAT?**

BAT is given as an injection into your vein. Your doctor will determine the dose of BAT. The treatment may take several hours to administer. Your doctor will decide if you need more than one injection.

### **What are the possible or reasonably likely side effects of BAT?**

The most common side effects of BAT are:

- Headache

- Fever
- Rash
- Hives
- Chills
- Nausea
- Swelling

Some people have a chilly feeling, difficulty breathing, and have a quick rise in body temperature within the first 20 to 60 minutes after getting BAT. This can be managed by your doctor.

BAT can cause allergic reactions. Tell your doctor or go to the emergency department right away if you have trouble breathing, swelling of your tongue or lips, or a very fast heart rate because this can be signs of a serious allergic reaction.

Tell your doctor if you get pains in your joints and back, fever, and a rash within one to three weeks after getting BAT. These can be signs of “serum sickness” and can last for a few weeks. Your doctor can give you medicine to help with serum sickness.

Talk to your doctor about any side effects that concern you. You can ask your doctor for additional prescribing information that is available to healthcare professionals.

**What other information do you need to know about BAT?**


BAT is made from horse plasma. The horses are carefully screened and the plasma is carefully cleaned, but there is a small risk that it may give you a virus. Talk to your doctor if you have any symptoms that concern you.

You may report side effects directly to Cangene Corporation at 1-800-768-2304 or to the FDA’s MedWatch reporting system at 1-800-FDA-1088.

**Manufactured By:**

Cangene Corporation  
Winnipeg, Manitoba  
Canada, R3T 5Y3


## Top Label reflective of IND Status



## Back of the top label



## Bottom Label reflective of future commercial status



**Licensed Shelf Carton Label (6 Vials)**

<b>STRATEGIC NATIONAL STOCKPILE USE ONLY</b>	Contains no preservatives Intravenous use only Store at $\leq -15^{\circ}\text{C}$ ( $\leq 5^{\circ}\text{F}$ ) See package insert for directions Rx Only		<b>(01) 003 6049200752 (BAR CODE)</b>
<b>Botulism Antitoxin Heptavalent (A, B, C, D, E, F, G) – (EQUINE)</b>			
<b>POTENCY:</b> Serotype A: > 4500 U/vial Serotype B: > 3300 U/vial Serotype C: > 3000 U/vial	Serotype D: > 600 U/vial Serotype E: > 5100 U/vial	Serotype F: > 3000 U/vial Serotype G: > 600 U/vial	
Lot Number:	Manufactured by: Cangene Corporation Winnipeg, Manitoba, Canada, R3T 5Y3 U.S. License No. 1201		
Fill Volume:			
<b>Expiry Date:</b>	Distributed by: Centers for Disease Control and Prevention, Atlanta, GA		
Number of vials: 6  3501xxxx			

**Licensed Shelf Carton Label (24 Vials)**

<b>STRATEGIC NATIONAL STOCKPILE USE ONLY</b>	Contains no preservatives Intravenous use only Store at $\leq -15^{\circ}\text{C}$ ( $\leq 5^{\circ}\text{F}$ ) See package insert for directions Rx Only	
<b>Botulism Antitoxin Heptavalent (A, B, C, D, E, F, G) – (EQUINE)</b>		
<b>POTENCY:</b> Serotype A: > 4500 U/vial Serotype B: > 3300 U/vial Serotype C: > 3000 U/vial	Serotype D: > 600 U/vial Serotype E: > 5100 U/vial	Serotype F: > 3000 U/vial Serotype G: > 600 U/vial
Lot Number:	Manufactured by:	
Fill Volume:	Cangene Corporation Winnipeg, Manitoba, Canada, R3T 5Y3	
<b>Expiry Date:</b>	U.S. License No. 1201	
Number of vials: 24	Distributed by:	
3501xxxx	Centers for Disease Control and Prevention, Atlanta, GA	
		(01) 003 6049200753 (BAR CODE)